

## GLP-1 AS AN ADJUNCT TO PROLACTIN AND ANTI-CD3 IN TYPE 1 DIABETES TREATMENT

Clement Chan, Vipul Shrivastava, Colin Hyslop & Carol Huang

Department of Biochemistry & Molecular Biology, Cumming School of Medicine, University of Calgary  
ckochan@ucalgary.ca

### INTRODUCTION

Type 1 diabetes mellitus (T1DM) is an autoimmune disease where the insulin-producing pancreatic  $\beta$ -cells are destroyed [1]. Affected individuals would have insulin deficiency, leading to the development of hyperglycemia, and unless insulin is provided, the patients will succumb to the disease [1]. The specific triggers of T1DM is still unknown, but many have hypothesized that hygiene and other environmental factors contribute to the pathogenesis of T1DM [1].

Anti-CD3 (aCD3) is a monoclonal antibody that is known to modulate immunity and stop the autoimmune attack on  $\beta$ -cells by T-cells [2]. Prolactin (PRL) is a peptide hormone that plays an important regulatory role in  $\beta$ -cell adaptation to pregnancy [3]. Its known effects include upregulating  $\beta$ -cell mass, proliferation, insulin secretion, and downregulating apoptosis [3,4]. Glucagon-like peptide 1 (GLP-1) is an incretin that is secreted by the L-cells of the distal ileum and exhibits effects such as increasing  $\beta$ -cell mass, proliferation, insulin secretion and satiety while decreasing glucagon secretion and  $\beta$ -cell apoptosis [5,6]. The objectives of this research are to determine if the addition of GLP-1 to PRL+aCD3 treatment would improve  $\beta$ -cell function in diabetic mice and to determine the source of the  $\beta$ -cells in the cured mice.

### METHODS

This experiment was conducted using a non-obese diabetic (NOD) mice model. After the T1DM onset in the NOD mice, aCD3 (10  $\mu$ g/day) was administered to the mice for 5 days, PRL (2.7  $\mu$ g/day) for 21 days, and GLP-1 (10  $\mu$ g/day) for 21 days. Upon reaching the 40<sup>th</sup> week after initial diabetes onset, the mice were sacrificed and the pancreases were taken for immunological studies.  $\beta$ -cell mass, neogenesis, proliferation, and apoptosis were determined by performing immunohistochemistry (IHC). Insulin ELISA was used as an assay for pancreatic insulin content and secretion.

### RESULTS

The GLP-1+PRL+aCD3 treatment group had a survival rate of 83.3% (n=6) resulting in a ~1.8-fold increase over the PRL+aCD3 group (n=11) and a ~3.3-fold increase over the aCD3 group (n=16).

	GLP-1 +PRL+aCD3	PRL+aCD3	aCD3
<b><math>\beta</math>-cell fraction</b>	0.37 $\pm$ 0.10%* (n=4)	0.41 $\pm$ 0.11%* (n=7)	0.09 $\pm$ 0.03% (n=7)
<b>RFP+ <math>\beta</math>-cell fraction</b>	91.48 $\pm$ 0.36% (n=2)	96.31 $\pm$ 1.52% (n=4)	93.51 $\pm$ 0.73% (n=2)
<b>BrdU+ <math>\beta</math>-cell fraction</b>	4.65 $\pm$ 1.77%* (n=3)	5.21 $\pm$ 3.30%* (n=7)	3.57 $\pm$ 2.00% (n=4)
<b>TUNEL+ <math>\beta</math>-cell fraction</b>	0.17 $\pm$ 0.09% (n=3)	0.54 $\pm$ 0.16% (n=2)	0.78 $\pm$ 0.27% (n=2)

**Figure 1.**  $\beta$ -cell fraction, RFP+, BrdU+, and TUNEL+  $\beta$ -cell fractions in various groups. “\*”: p<0.05 in comparison to the aCD3 group. “n” = number of mice in each group.

### DISCUSSION AND CONCLUSIONS

The group treated with GLP-1+PRL+aCD3 had comparable  $\beta$ -cell mass to the PRL+aCD3 group. Evidence for  $\beta$ -cell neogenesis was determined by quantifying RFP+  $\beta$ -cells. When RFP is expressed on a  $\beta$ -cell, it indicates that the  $\beta$ -cell is pre-existing and is not from neogenesis. Neogenesis did not contribute to the  $\beta$ -cell mass since all three groups shared a similar RFP+ fraction at a value that is near 100%. BrdU was incorporated in the mice’s DNA to act as a cell proliferation marker. The BrdU+  $\beta$ -cell fractions were similar between the GLP-1+PRL+aCD3 and PRL+aCD3 groups, suggesting that GLP-1 does not stimulate  $\beta$ -cells proliferation, above and beyond the effect of PRL. GLP-1, however, may decrease the apoptosis of  $\beta$ -cells, as suggested by the lower TUNEL+  $\beta$ -cell fraction in the GLP-1+PRL+aCD3 treatment group when compared with the PRL+aCD3 group.

GLP-1 was determined to increase the survival of NOD mice, however, its underlying mechanisms is still unknown. Determining pancreatic insulin content and GLUT2 expression in future experiments would identify such mechanisms.

### REFERENCES

1. Daneman D. *Lancet*. **358**:847-858,2006.
2. Herold KC & Taylor L. *Immunol Res*. **28**:141-150, 2003.
3. Huang C. *J Physio*. **591**:1325-1338, 2013.
4. Nielsen JH, et al. *J Mol Med*. **77**:62-66, 1999.
5. Brubaker PL & Drucker DJ. *Endocrinology*. **145**:2653-2659, 2004.
6. Sherry NA, et al. *Endocrinology*. **148**:5136-5144, 2007.